



1. Consider the points $A = (2, 0, 1)$ and $B = (4, 2, 5)$ in \mathbb{R}^3 .

(a) Find the point M which is halfway between A and B on the line segment L joining them. (2 pts)

$$M = \vec{v} + \frac{1}{2}(\vec{w} - \vec{v}) = \frac{1}{2}(\vec{v} + \vec{w}) = \frac{1}{2}((2, 0, 1) + (4, 2, 5)) \\ = \frac{1}{2}(6, 2, 6) = (3, 1, 3)$$



(b) Find the equation for the plane P consisting of all points that are equidistant from A and B . (3 pts)



Point on P : $M = (3, 1, 3)$

Normal to P : $\frac{1}{2}(\vec{w} - \vec{v}) = \frac{1}{2}((4, 2, 5) - (2, 0, 1)) \\ = \frac{1}{2}(2, 2, 4) = (1, 1, 2)$

Ans: $1 \cdot (x-3) + 1 \cdot (y-1) + 2 \cdot (z-3) = 0$

2. Consider the function

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

(a) Compute the following limit, if it exists. (4 pts)

$$\lim_{(x, y) \rightarrow (0, 0)} f(x, y)$$

Along x -axis: $f(x, 0) = \frac{x \cdot 0}{x^2 + 0^2} = 0$

Along $y = x$: $f(x, x) = \frac{x \cdot x}{x^2 + x^2} = \frac{1}{2}$













So as f does not approach a consist. value as $(x, y) \rightarrow (0, 0)$
the limit does not exist

(b) Where on \mathbb{R}^2 is the function f continuous? (1 pts)

Everywhere except $(0, 0)$.

3. Match the following functions $\mathbb{R}^2 \rightarrow \mathbb{R}$ with their graphs and contour diagrams. Here each contour diagram consists of level sets $\{f(x, y) = c_i\}$ drawn for evenly spaced c_i . (9 pts)

- (a) $\sqrt{8 - 2x^2 - y^2}$ (b) $\cos x$ (c) xy



4. Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $f(x, y) = xy$.

(a) Use Lagrange multipliers to find the global (absolute) max and min of f on the circle

$$x^2 + y^2 = 2. \quad (6 \text{ pts})$$

First $\underbrace{g(x, y)}$

$$\nabla f = \nabla g = \lambda \nabla g = \lambda(2x, 2y).$$

Thus $y = 2\lambda x$ and $x = 2\lambda y$ and so $2\lambda = \frac{x}{y} = \frac{y}{x}$.

Therefore $x^2 = y^2$. Since also $x^2 + y^2 = 2$, we get $2x^2 = 2$

$\Rightarrow x^2 = \pm 1$ and $y^2 = \pm 1$. So there are 4 critical pts

Point	(1, 1)	(1, -1)	(-1, 1)	(-1, -1)
Val of f	1	-1	-1	+1
Type	Max	Min	Min	Max

Note: Global min/max exist since C is closed and bounded.

(b) If they exist, find the global min and max of f on $D = \{x^2 + y^2 \leq 2\}$. (2 pts)

Need to also check for crit pts $\nabla f = 0$ inside D .

As $\nabla f = (2x, 2y)$, only such pt is $(0, 0)$ where $f = 0$.

So global min/max are the same as in (a).

(c) For each critical point in the interior of D you found in part (b), classify it as a local min, local max, or saddle. (2 pt)

Check

$$D = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1 < 0 \quad \text{hence a saddle.}$$

(d) If they exist, find the global min and max of f on \mathbb{R}^2 . (2 pts)

Neither exist since $f \rightarrow +\infty$ along $y = x$ and $\rightarrow -\infty$ along $y = -x$.

5. A function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ takes on the values shown in the table at right.

(a) Estimate the partials $f_x(1, 1)$ and $f_y(1, 1)$. (2 pts)

$$f_x(1, 1) \approx \frac{f(1.4, 1) - f(1, 1)}{0.4} = \frac{3.4 - 3.0}{0.4} = 1$$

$$f_y(1, 1) \approx \frac{f(1, 1.4) - f(1, 1)}{0.4} = \frac{3.8 - 3.0}{0.4} = 2$$

	x				
	0.2	0.6	1.0	1.4	1.8
1.8	3.16	3.88	4.60	5.32	6.04
1.4	2.68	3.24	3.80	4.36	4.92
1.0	2.20	2.60	3.00	3.40	3.80
0.6	1.72	1.96	2.20	2.44	2.68
0.2	1.24	1.32	1.40	1.48	1.56

(b) Use your answer in (a) to approximate $f(1.1, 1.2)$. (2 pts)

$$f(1.1, 1.2) \approx f(1, 1) + f_x(1, 1)(\Delta x) + f_y(1, 1)(\Delta y)$$

$$\approx 3 + 1(0.1) + 2(0.2) = 3.5$$


(c) Determine the sign of $f_{xy}(1, 1)$: positive negative zero (1 pt)

$$f_y(1.4, 1) \approx \frac{f(1.4, 1.4) - f(1.4, 1)}{0.4} = \frac{4.36 - 3.40}{0.4} = \frac{0.96}{0.4} > 2$$

Thus f_y increases as x increases $\Rightarrow \frac{\partial}{\partial x} f_y > 0$.


6. Consider the region E shown at right, which is bounded by the xy -plane, the plane $z - y = 0$ and the surface $x^2 + y = 1$. Complete setup, but do not evaluate, a triple integral that computes the volume of E . (6 pts)

$$\int_{-1}^1 \int_0^{1-x^2} \int_0^y 1 \, dz \, dy \, dx$$



7. Consider the portion R of the cylinder $x^2 + y^2 \leq 2$ which lies in the positive octant and below the plane $z = 1$. Compute the total mass of R when it is composed of material of density $\rho = e^{x^2+y^2}$.

(7 pts)



$$\text{Mass} = \iiint_R \rho \, dV = \iiint_R e^{x^2+y^2} \, dV$$

$$= \int_0^{\sqrt{2}} \int_0^1 \int_0^{\pi/2} e^{r^2} r \, d\theta \, dz \, dr = \int_0^{\sqrt{2}} \frac{\pi e^{r^2} r \, dr}{2}$$

$u=r^2$
 $du=2r \, dr$

using
cylindrical
coord.

$$= \int_0^2 \frac{\pi e^u}{4} \, du$$

$$= \frac{\pi}{4} (e^2 - 1).$$

8. For the curve C in \mathbb{R}^2 shown and the vector field $F = (\ln(\sin(x)), \cos(\sin(y)) + x)$ evaluate

$\int_C F \cdot d\vec{r}$ using the method of your choice. (5 pts)


Let's use Green's Thm:

$$\int_C \vec{F} \cdot d\vec{r} = - \iint_D \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \, dA$$


because
 C is clockwise

$$= - \iint_D 1 - 0 \, dA$$

$$= - \text{Area}(\triangle) = -\frac{1}{2}$$



9. Let R be the region shown at right.



(a) Find a transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ taking $S = [-1, 1] \times [-1, 1]$ to R . (4 pts)

Want $T(1, 0) = (1, 1)$ $T(0, 1) = (-1, 1)$. If

$T(u, v) = (au + bv, cu + dv)$ we can solve $T(1, 0) = (a, c) = (1, 1)$
 $T(0, 1) = (b, d) = (-1, 1)$
 to find $T(u, v) = (u - v, u + v)$.

Check: $T(1, 1) = (1 - 1, 1 + 1) = (0, 2)$
 $T(1, -1) = (2, 0)$

(b) Use your change of coordinates to evaluate $\iint_R y^2 dA$ via an integral over S . (6 pts)

Emergency backup transformation: If you can't do (a), pretend you got the answer $T(u, v) = (uv, u + v)$ and do part (b) anyway.

$$\iint_R y^2 dA = \iint_S (u+v)^2 |\det J| du dv$$

$$= \int_{-1}^1 \int_{-1}^1 2(u^2 + 2uv + v^2) du dv$$

$$= 2 \int_{-1}^1 \left. \frac{u^3}{3} + u^2 v + v^2 u \right|_{u=-1}^1 dv$$

$$= 2 \int_{-1}^1 \left(\frac{2}{3} + 2v^2 \right) dv = 4 \left(\frac{1}{3} v + \frac{v^3}{3} \right) \Big|_{v=-1}^1$$

$$= \frac{16}{3}$$

$T = (g, h)$





$$J = \begin{pmatrix} g_u & g_v \\ h_u & h_v \end{pmatrix}$$

$$= \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$$

$\det J = 2$

10. Consider the surface S which is parameterized by $\mathbf{r}(u, v) = (\sqrt{1+u^2} \cos v, \sqrt{1+u^2} \sin v, u)$ for $-1 \leq u \leq 1$ and $0 \leq v \leq 2\pi$.

(a) Circle the picture of S . (2 pts)



(b) Completely setup, but do not evaluate, an integral that computes the surface area of S . (6 pts)

$$\begin{aligned} \text{Area} &= \iint_S 1 \, dA = \iint_D |\vec{r}_u \times \vec{r}_v| \, du \, dv \\ &= \int_0^{2\pi} \int_{-1}^1 \sqrt{1+2u^2} \, du \, dv \end{aligned}$$


$$\begin{aligned} \vec{r}_u \times \vec{r}_v &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{u}{\sqrt{1+u^2}} \cos v & \frac{u}{\sqrt{1+u^2}} \sin v & 1 \\ -\sqrt{1+u^2} \sin v & \sqrt{1+u^2} \cos v & 0 \end{vmatrix} \\ &= (\sqrt{1+u^2} \cos v, -\sqrt{1+u^2} \sin v, u) \end{aligned}$$

$$\begin{aligned} |\vec{r}_u \times \vec{r}_v| &= \sqrt{(1+u^2) \cos^2 v + (1+u^2) \sin^2 v + u^2} \\ &= \sqrt{1+2u^2} \end{aligned}$$

11. For the cone S at right, give a parameterization $\mathbf{r}: D \rightarrow S$. Explicitly specify the domain D . (5 pts)

Params: $u = y$ $v = \text{angle about } y\text{-axis}$

Radius about y -axis is a fn of $y = u$




with:

$$\vec{r}(u, v) = ((1-u) \cos v, (1-u) \sin v, u)$$





So take D

$$= \{0 \leq u \leq 1 \text{ and } 0 \leq v \leq 2\pi\}$$



12. Consider the region R in \mathbb{R}^3 above the surface $x^2 + y^2 - z = 4$ and below the xy -plane. Also consider the vector field $F = (0, 0, z)$.

(a) Circle the picture of R below. (2 pts)



(b) Directly calculate the flux of F through the entire surface ∂R , with respect to the outward unit normals. (10 pts)

Now $\partial R = (T = \text{disk})$ and $(S = \text{bowl})$. First, $\iint_T \vec{F} \cdot \vec{n} dA = \iint_T (0, 0, 0) \cdot (0, 0, 1) dA = \iint_T 0 dA = 0$. Second, let's param. S by $\vec{r}(u, v) = (u, v, u^2 + v^2 - 4)$ on $D = \{u^2 + v^2 \leq 4\}$. Then $\vec{r}_u \times \vec{r}_v = \begin{vmatrix} i & j & k \\ 1 & 0 & 2u \\ 0 & 1 & 2v \end{vmatrix} = (-2u, -2v, 1)$. As this points the wrong way we use $\vec{r}_v \times \vec{r}_u$ instead. Now Flux = $\iint_S (\vec{F} \cdot \vec{n}) dA = \iint_D \vec{F}(\vec{r}(u, v)) \cdot (\vec{r}_v \times \vec{r}_u) du dv = \iint_D (0, 0, u^2 + v^2 - 4) \cdot (2u, 2v, -1) du dv = \iint_D 4 - u^2 - v^2 du dv = \int_0^2 \int_0^{2\pi} (4 - r^2) r d\theta dr = 2\pi \int_0^2 4r - r^3 dr = 2\pi \left(2r^2 - \frac{r^4}{4} \right) \Big|_{r=0}^{r=2} = 2\pi (8 - 4) = \boxed{8\pi}$

Hence $\iint_{\partial R} \vec{F} \cdot \vec{n} dA = \iint_T \vec{F} \cdot \vec{n} dA + \iint_S \vec{F} \cdot \vec{n} dA = 0 + 8\pi = \boxed{8\pi}$

(c) Use the Divergence Theorem and your answer in (b) to compute the volume of R . (3 pts)


$$\iint_{\partial R} \vec{F} \cdot \vec{n} dA = \iiint_R \operatorname{div} \vec{F} dV = \iiint_R 1 dV = \text{Volume.}$$

$$\frac{\partial 0}{\partial x} + \frac{\partial 0}{\partial y} + \frac{\partial z}{\partial z} = 1$$

So: $\boxed{\text{Vol} = 8\pi}$

13. Let C be the curve shown at right, which is the boundary of the portion of the surface $x + z^2 = 1$ in the positive octant where additionally $y \leq 1$.

- (a) Label the four corners of C with their (x, y, z) -coordinates. (1 pt)



- (b) For $F = (0, xyz, xyz)$, directly compute $\int_C F \cdot dr$. (6 pts)

Break C up into C_1, C_2, C_3, C_4

Notice if any of x, y or z is 0, then $\vec{F} = \vec{0}$. Thus for any C_i except C_1 , we have $\int_{C_i} \vec{F} \cdot d\vec{r} = \int_{C_i} \vec{0} \cdot d\vec{r} = 0$.

For C_1 , we parameterize $-C_1$ via

$$\vec{r}(t) = (1 - t^2, 1, t) \text{ for } 0 \leq t \leq 1.$$

$$\begin{aligned} \text{So } \int_{C_1} \vec{F} \cdot d\vec{r} &= - \int_{-C_1} \vec{F} \cdot d\vec{r} = - \int_0^1 (0, t - t^3, t - t^3) \cdot (-2t, 0, 1) dt \\ &= \int_0^1 t^3 - t dt = \left. \frac{t^4}{4} - \frac{t^2}{2} \right|_{t=0}^1 = \frac{1}{4} \end{aligned}$$

- (c) Compute $\text{curl } F$. (2 pts) Hence $\int_C \vec{F} \cdot d\vec{r} = \sum \int_{C_i} \vec{F} \cdot d\vec{r} = 0 + 0 + 0 + \frac{1}{4}$

$$\text{curl } \vec{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 0 & xyz & xyz \end{vmatrix} = (xz - xy, -yz, yz)$$

- (d) Use Stokes' Theorem to compute the flux of $\text{curl } F$ through the surface S where the normals point out from the origin. (3 pts)

$$\iint_S (\text{curl } \vec{F}) \cdot \vec{n} dA = \int_C \vec{F} \cdot d\vec{r} = \frac{1}{4}$$

since orient of C doesn't mesh with \vec{n} .

- (e) Give two distinct reasons why the vector field F is not conservative. (2 pts)

$$\text{curl } \vec{F} \neq 0 \text{ and } C \text{ is a closed curve with } \int_C \vec{F} \cdot d\vec{r} \neq 0.$$