

Lecture 23: Whitehead's Theorem

Whitehead's Thm: $f: X \rightarrow Y$ a map between ^{connected} CW complexes.

If f_* is an \cong on all π_n then f is a homotopy equivalence. Moreover, f is the inclusion of a subcomplex, then Y deformation retracts to X .

Compression Lemma: (X, A) a CW pair, (Y, B) any pair $\neq \emptyset$.

For each n where $X \setminus A$ has n -cell, assume $\pi_n(Y, B, y_0) = 0$ for all $y_0 \in B$. Then every map $f: (X, A) \rightarrow (Y, B)$ is homotopic, rel A , to a map $f: X \rightarrow B$.

Pf of Lemma: Inductively, build $f_k: (X, A) \rightarrow (Y, B)$

so that ① f_k is homotopic to f ; rel A .


② $f_k(X^k \cup A) \subseteq B$. ③ homotopy from f_{k-1} to f_k is const on $X^{k-1} \cup A$.

[f_{k+1} build from f_k by first changing on X^k , then applying the homotopy extension thm.]

Now define $f_\infty: (X, A) \rightarrow (Y, B)$ by

If $f: X \rightarrow Y$ is any map, consider the

mapping cylinder $M_f = \frac{X \times [0,1] \amalg Y}{(x,1) \sim f(x)}$



Note: M_f def retracts to Y ; $X \hookrightarrow M_f \xrightarrow[\cong]{\text{retract}} Y$
 $\underbrace{\hspace{10em}}_f$

Claim: When f_* is an \cong on π_n , then M_f deformation retracts to X . [$\Rightarrow X \hookrightarrow M_f$ is a homotopy equiv, hence $X \xrightarrow{f} Y$ is.]


If f is cellular, i.e. $f(X^k) \subseteq Y^k$ for all k , then

M_f is a CW complex

with $X \times \{0\}$ as a subcomplex,

allowing us to apply the

earlier case. So have reduced Whitehead's Thm to:



Thm: Every $f: X \rightarrow Y$ of CW complexes is homotopic to a cellular map. If f is already cellular on a subcomplex A of X , then the homotopy is const on A .


Cor: $\pi_n(S^k) = 0$ for $n < k$.

$$f_\infty|_{X^k} = f_k|_{X^k} = f_j|_{X^k} \text{ for } j \geq k. \quad (2)$$

Then f_∞ is homotopic to f via a homotopy that implements the homotopy from f_k to f_{k+1} in the time interval $[1-2^{-k}, 1-2^{-(k+1)}]$. \square

Works since X has the weak topology: $U \subseteq X$ is open $\Leftrightarrow U \cap X^k$ is open for each k .

Pf of Thm: Suppose X is a subcomplex and f is inclusion

Long exact sequence gives (for any $x_0 \in X$) 

$$\begin{array}{ccccccc} \rightarrow \pi_n X & \xrightarrow{f_*} & \pi_n Y & \rightarrow & \pi_n(Y, X) & \rightarrow & \pi_{n-1}(X) \xrightarrow{\cong} \pi_{n-1}(Y) \rightarrow \\ & \cong & \searrow \circ & \nearrow & \searrow \circ & \nearrow & \cong \end{array}$$


and so $\pi_n(Y, X) = 0$. Applying the Compression Lemma to $\text{id}: (Y, X) \rightarrow (Y, X)$ gives the needed deformation retraction

Pf of Cor: Consider the usual cell decomp of S^n (4) and S^k with one zero cell and one other cell. (the base pt)

Then the Thm implies that any map $S^n \rightarrow S^k$ is homotopic (rel base pt) to the const map. \square

[Query: how did you show $\pi_1(S^2) = 1$?]


Pf of Cellular Approximation: Idea



Proceed inductively: (1) Homotope $f|_{X^0}$ so that $f(X^0) \subseteq Y^0$; extend to all of X by homotopy extension thm.

(2) Homotope $f: X^1 \rightarrow Y$, rel X^0 , so that $f(X^1) \subseteq Y^1$ by picking p_i in each cell of $Y \setminus Y^1$ not in image and pushing outwards.

Again extend to all of X by homotopy ext. thm.



(n) Repeat for each n in the same manner.

(5)

[Query: What is missing here?]

Lemma: $Z = W \cup$ ^{some σ} $(k\text{-cell } e^k)$. For $n < k$,
and map $f: I^n \rightarrow Z$ is homotopic, rel $f^{-1}(W)$,
to a map where $g(I^n)$ is a proper subset
of $\text{int}(e^k)$.

Lemma: Any cpt A in a CW complex X
meets the interiors of finitely many cells.