

# Lecture 5: Using matrices to solve linear systems (1)

Reference: [SRREF of Breezer]

Last time: System of linear equations:

Variables:  $x_1, x_2, \dots, x_n$

Equations:

$$\begin{aligned} a_{11}x_1 + \dots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + \dots + a_{2n}x_n &= b_2 \\ \vdots & \\ a_{m1}x_1 + \dots + a_{mn}x_n &= b_m \end{aligned}$$

where  $a_{ij}$  and  $b_i$  are in  $\mathbb{R}$ .

---

To a linear system, we associate a coefficient matrix  $A = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \dots & a_{mn} \end{pmatrix}$ ,

a vector of constants  $b = \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix}$  ← column vector.

and an augmented matrix

$$\begin{pmatrix} a_{11} & \dots & a_{1n} & b_1 \\ \vdots & & \vdots & \vdots \\ a_{m1} & \dots & a_{mn} & b_m \end{pmatrix}$$

$$\underline{\text{Ex:}} \quad x_1 + 2x_2 + 2x_3 = 4$$

$$x_1 + 3x_2 + 3x_3 = 5$$

$$2x_1 + 6x_2 + 5x_3 = 6$$

(2)

This system has  $A = \begin{pmatrix} 1 & 2 & 2 \\ 1 & 3 & 3 \\ 2 & 6 & 5 \end{pmatrix}$  and  $b = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}$ .

The augmented matrix  $(A|b)$  is  $\begin{pmatrix} 1 & 2 & 2 & 4 \\ 1 & 3 & 3 & 5 \\ 2 & 6 & 5 & 6 \end{pmatrix}$ .

Decoding:

Given an  $m \times n$  matrix  $A$  and an  $m$  entry column vector  $b$ , we associate a linear system  $\mathcal{L}(A, b)$ . Also works for augmented matrices.

$$\underline{\text{Ex:}} \quad \mathcal{L}\left(\begin{pmatrix} 1 & 3 & 1 \\ 2 & 1 & 0 \end{pmatrix}\right) = \begin{cases} x_1 + 3x_2 = 1 \\ 2x_1 + x_2 = 0 \end{cases}$$

Row operations: These take a matrix  $M$  and give a new matrix of the same size.

- ① Swap two rows of  $M$
- ② Multiply all entries in a row by some nonzero  $c$  in  $\mathbb{R}$
- ③ Add a multiple of one row to another.

[Hint at use for solving lin. systems.]

Ex:  $M = \begin{pmatrix} 1 & 2 & 2 & 4 \\ 1 & 3 & 3 & 5 \\ 2 & 6 & 5 & 6 \end{pmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{pmatrix} 1 & 2 & 2 & 4 \\ 2 & 6 & 5 & 6 \\ 1 & 3 & 3 & 5 \end{pmatrix}$

$\xrightarrow{-3R_3} \begin{pmatrix} 1 & 2 & 2 & 4 \\ 2 & 6 & 5 & 6 \\ -3 & -9 & -9 & -15 \end{pmatrix} \xrightarrow{3R_1 + R_3} \begin{pmatrix} 1 & 2 & 2 & 4 \\ 2 & 6 & 5 & 6 \\ 0 & -3 & -3 & -3 \end{pmatrix}$

$\xrightarrow{-\frac{1}{3}R_3} \begin{pmatrix} 1 & 2 & 2 & 4 \\ 2 & 6 & 5 & 6 \\ 0 & 1 & 1 & 1 \end{pmatrix}$

Def: A matrix  $M$  is row equivalent to  $N$  if there is a sequence of row operations that turns  $M$  into  $N$ .

Note: On HW you'll show this is symmetric in  $M$  and  $N$ .

If numbers  $(x_1, \dots, x_n)$  satisfy  $(E_1)$  (5)

then they also sat  $(E_2)$ . Conversely if

they sat  $(E_2)$  they also sat  $(E_1)$  since we can subtract the first eqn from the second equation of  $(E_2)$ .

So all three kinds of row ops don't change the sol set of the corresp linear sys.  $\square$

Ex. (Continuation)

$$M \xrightarrow[\text{ops}]{\text{row}} \begin{pmatrix} 1 & 2 & 2 & 4 \\ 2 & 6 & 5 & 6 \\ 0 & 1 & 1 & 1 \end{pmatrix} \xrightarrow{-2R_1 + R_2} \begin{pmatrix} 1 & 2 & 2 & 4 \\ 0 & 2 & 1 & -2 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

$$\xrightarrow{-R_2 + R_1} \begin{pmatrix} 1 & 0 & 1 & 6 \\ 0 & 2 & 1 & -2 \\ 0 & 1 & 1 & 1 \end{pmatrix} \xrightarrow{-2R_3 + R_2} \begin{pmatrix} 1 & 0 & 1 & 6 \\ 0 & 0 & -1 & -4 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

$$\xrightarrow[\begin{matrix} R_2 + R_1 \\ R_2 + R_3 \end{matrix}]{R_2 + R_1} \begin{pmatrix} 1 & 0 & 0 & 2 \\ 0 & 0 & -1 & -4 \\ 0 & 1 & 0 & -3 \end{pmatrix} \xrightarrow{-R_2} \begin{pmatrix} 1 & 0 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 1 & 0 & -3 \end{pmatrix}$$

$$\xrightarrow{\text{ZS}} \begin{matrix} x_1 = 2 \\ x_3 = 4 \\ x_2 = -3 \end{matrix} \quad \text{So } (x_1, x_2, x_3) = (2, -3, 4)$$

Thm: If  $M$  is row equivalent to  $N$ , then the linear systems  $LS(M)$  and  $LS(N)$  have the same solution sets.

Pf: Just need to check that a single row op does not change the solution set of the corresp. linear system. Row op ① is fine. [since it corresponds to reordering two equations]

Row op ② just scales the assoc. eqn. For row op ③, it is enough to consider the case of adding rows without any scaling.

[The general case follows by using row op ②]

On the equation side, have

①  $a_1x_1 + \dots + a_nx_n = b$   
 $a'_1x_1 + \dots + a'_nx_n = b'$   $\rightsquigarrow$

②  $a_1x_1 + \dots + a_nx_n = b$   
 $(a'_1 + a_1)x_1 + \dots + (a'_n + a_n)x_n = b + b'$

is the unique soln to our original system: ⑥

$$x_1 + 2x_2 + 2x_3 = 4$$

$$x_1 + 3x_2 + 3x_3 = 5$$

$$2x_1 + 6x_2 + 5x_3 = 6.$$

as you can (and should!) double check:

Goal: Develop systematic procedure for solving such systems, by finding matrices like these:

Reduced Row-Echelon Form: A matrix where:

- ① All zero rows are at the bottom.
- ② The leftmost entry of every row is 1, and is called a leading 1.
- ③ A leading 1 is the only nonzero entry in its column.
- ④ Suppose entries  $(i, j)$  and  $(s, t)$  are leading 1's. If  $s > i$  then  $t > j$ .