

## Math 416: HW 2 due Friday, February 2, 2018.

Webpage: <http://dunfield.info/416>

Office hours: Tue 10–11, Wed 3:30–4:30, Thur 2–3 and by appointment. My office is 378 Altgeld.

Textbooks: In the assignment, the two texts are abbreviated as follows:

[FIS] Freidberg, Insel, Spence, *Linear Algebra*, 4th edition, 2002.

[B] Breezer, *A First Course in Linear Algebra*, Version 3.5, 2015.

### Problems:

- In these questions, you will determine whether one vector is a linear combination of two others.
  - Section 1.4 of [FIS], Problem 3: parts (a) and (c).
  - Section 1.4 of [FIS], Problem 4: parts (a) and (e).
- In class, I defined the span of a finite list of vectors  $u_1, u_2, \dots, u_n$ . More generally, given a nonempty subset  $S$  of a vector space  $V$ , one defines  $\text{span}(S)$  to be the set of all linear combinations of vectors in  $S$ . Here are some problems about the span.
  - Section 1.4 of [FIS], Problem 5: parts (g) and (h).
  - Suppose  $S_1$  and  $S_2$  are subsets of a vector space  $V$ . Show that if  $S_1$  is contained in  $S_2$ , then  $\text{span}(S_1)$  is contained in  $\text{span}(S_2)$ .
  - Let  $V = \mathbb{R}^2$  and  $S = \{(x, y) \text{ where } x \geq 0 \text{ and } y \geq x\}$ . Find  $\text{span}(S)$ .
- Solve each of the following linear systems by writing down its augmented matrix, doing row operations to get a matrix in reduced row echelon form, and using that to find all of the solutions. You should label your row operations as in §RREF of [B].
  -

$$2x_1 + x_2 = 0$$

$$x_1 + x_2 = 1$$

$$3x_1 + 4x_2 = 5$$

$$3x_1 + 5x_2 = 7$$

(b)

$$\begin{aligned}y_1 + 2y_2 - y_3 &= 1 \\y_1 + y_2 + 2y_3 &= 0 \\5y_1 + 8y_2 + y_3 &= 1\end{aligned}$$

(c)

$$\begin{aligned}2x_1 + 4x_2 + 5x_3 + 7x_4 &= 18 \\x_1 + 2x_2 + x_3 - x_4 &= 3 \\4x_1 + 8x_2 + 7x_3 + 5x_4 &= 24\end{aligned}$$

4. Suppose that  $A$ ,  $B$ , and  $C$ , are  $m \times n$  matrices with real coefficients. Prove the following three facts from the definition of row equivalence.

(a)  $A$  is row equivalent to  $A$ .

(b) If  $A$  is row equivalent to  $B$ , then  $B$  is row equivalent to  $A$ .

(c) If  $A$  is row equivalent to  $B$ , and  $B$  is row equivalent to  $C$ , then  $A$  is row equivalent to  $C$ .

Note: A relationship that satisfies these three properties is known as an *equivalence relation*; this is a formal way of saying that a relationship behaves like equality, without requiring the relationship to be as strict as equality itself.

5. Suppose  $A$  is an  $m \times n$  matrix with real entries. The *null space* of  $A$ , denoted  $\mathcal{N}(A)$ , is the set of all solutions in  $\mathbb{R}^n$  to the linear system  $\mathcal{LS}(A, 0)$ , where here  $0$  is the zero vector in  $\mathbb{R}^m$ . Prove that  $\mathcal{N}(A)$  is a subspace of  $\mathbb{R}^n$ .