

Math 416: HW 2 due Friday, September 9, 2022.

Webpage: <http://dunfield.info/416>

Office hours: Wednesday 2:30–3:30pm and Thursday 3:00–4:00pm; other times possible by appointment. My office is 378 Altgeld.

Textbooks: In the assignment, the two texts are abbreviated as follows:

[FIS] Freidberg, Insel, Spence, *Linear Algebra*, 4th or 5th edition, 2002 or 2019.

[B] Breezer, *A First Course in Linear Algebra*, Version 3.5, 2015.

Problems:

- In these questions, you will determine whether one vector is a linear combination of two others.
 - Section 1.4 of [FIS], Problem 3: parts (a) and (c).
 - Section 1.4 of [FIS], Problem 4: parts (a) and (e).
- In class, I defined the span of a finite list of vectors u_1, u_2, \dots, u_n . More generally, given a nonempty subset S of a vector space V , one defines $\text{span}(S)$ to be the set of all linear combinations of vectors in S . Here are some problems about the span.
 - Section 1.4 of [FIS], Problem 5: parts (g) and (h).
 - Suppose S_1 and S_2 are subsets of a vector space V . Show that if S_1 is contained in S_2 , then $\text{span}(S_1)$ is contained in $\text{span}(S_2)$.
 - Let $V = \mathbb{R}^2$ and $S = \{(x, y) \text{ where } x \geq 0 \text{ and } y \geq x\}$. Find $\text{span}(S)$.
- Solve each of the following linear systems by writing down its augmented matrix, doing row operations to get a matrix in reduced row echelon form, and using that to find all of the solutions. You should label your row operations as in §RREF of [B].
 -

$$2x_1 + x_2 = 0$$

$$x_1 + x_2 = 1$$

$$3x_1 + 4x_2 = 5$$

$$3x_1 + 5x_2 = 7$$

(b)

$$y_1 + 2y_2 - y_3 = 1$$

$$y_1 + y_2 + 2y_3 = 0$$

$$5y_1 + 8y_2 + y_3 = 1$$

(c)

$$2x_1 + 4x_2 + 5x_3 + 7x_4 = 18$$

$$x_1 + 2x_2 + x_3 - x_4 = 3$$

$$4x_1 + 8x_2 + 7x_3 + 5x_4 = 24$$

4. Suppose that A , B , and C , are $m \times n$ matrices with real coefficients. Prove the following three facts from the definition of row equivalence.

(a) A is row equivalent to A .

(b) If A is row equivalent to B , then B is row equivalent to A .

(c) If A is row equivalent to B , and B is row equivalent to C , then A is row equivalent to C .

Note: A relationship that satisfies these three properties is known as an *equivalence relation*; this is a formal way of saying that a relationship behaves like equality, without requiring the relationship to be as strict as equality itself.

5. Suppose A is an $m \times n$ matrix with real entries. The *null space* of A , denoted $\mathcal{N}(A)$, is the set of all solutions in \mathbb{R}^n to the linear system $\mathcal{LS}(A, 0)$, where here 0 is the zero vector in \mathbb{R}^m . Prove that $\mathcal{N}(A)$ is a subspace of \mathbb{R}^n .