

Lecture 28: Proof of the diagonalization criteria. ①

Last time: λ eigenvalue of $A \in M_{n \times n}(\mathbb{R})$.

Multiplicities:

Algebraic: # of times $(t - \lambda)$ divides the char poly of A

Geometric: $\dim E_\lambda$

Thm: $A \in M_{n \times n}(\mathbb{R})$ is diagonalizable if and only if

a) The char poly of A splits completely over \mathbb{R}

b) (alg. mult) = (geom mult) for all eigenvalues of A .


_____ o _____

Lemma: Suppose $v_1, v_2, \dots, v_k \in \mathbb{R}^n$ are eigenvectors of A corresponding to distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$. Then $\{v_1, \dots, v_k\}$ are linearly independent.


Moral: Can't create an eigenvector with eigenvalue λ from eigenvectors with other eigenvalues.

Ex: $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$

$v_1 = e_1, \quad v_2 = e_2$



LA



$A(e_1 + e_2) = Ae_1 + Ae_2 = 2 \cdot e_1 + 3 \cdot e_2$

Proof of Lemma: Can assume $\lambda_k \neq 0$ for $k > 1$. Induct on k .

Base case: As v_1 is an eigenvector, it is nonzero and so $\{v_1\}$ is linearly independent.

Inductive Step: Assume $\{v_1, v_2, \dots, v_{k-1}\}$ is linearly independent. Will prove by contradiction, so assume $\{v_1, \dots, v_k\}$ is linearly dependent.

Then $v_k = a_1 v_1 + \dots + a_{k-1} v_{k-1}$ for some $a_i \in \mathbb{R}$.

Now
$$Av_k = \lambda_k v_k = \sum_{i=1}^{k-1} (\lambda_k a_i) v_i$$

and

$$Av_k = \sum_{i=1}^{k-1} A(a_i v_i) = \sum_{i=1}^{k-1} (\lambda_i a_i) v_i$$

Now $Av_k \neq 0$ but we have two distinct ③
ways of writing it as a linear combination
of the linearly indep. set $\{v_1, \dots, v_{k-1}\}$,
which is impossible. So $\{v_1, \dots, v_k\}$ is linearly
independent, completing the induction. \square

Lemma: Suppose $\lambda_1, \dots, \lambda_k$ are eigenvalues for A .
If $\beta_i \subseteq E_{\lambda_i}$ is lin. indep., then $\beta = \beta_1 \cup \dots \cup \beta_k$
is linearly independent.

Note: $\beta_i \cap \beta_j = \emptyset$ for $i \neq j$ as if $v \neq 0$ then
 $Av = \lambda_i v$ and $Av = \lambda_j v \Rightarrow \lambda_i = \lambda_j$.

Proof: Suppose $\beta_i = \{v_1^i, v_2^i, \dots, v_{d_i}^i\}$ and
there are scalars a_j^i such that

$$\sum_{i=1}^k \underbrace{\sum_{j=1}^{d_i} a_j^i v_j^i}_{w_i} = 0$$

Each $w_i \in E_{\lambda_i}$ and is either an eigenvector

for λ_i or is 0. By earlier lemma, can't ④

have a linear dependence among eigenvectors with different eigenvalues, so all $w_i = 0$. As each β_i is linearly indep, conclude $a_j^i = 0$ for all i and j .

So β is linearly independent. ▣

Proof of Thm: (\Leftarrow) By (a) have

$$\text{char poly of } A = \pm (t - \lambda_1)^{m_1} (t - \lambda_2)^{m_2} \cdots (t - \lambda_k)^{m_k}$$

for distinct $\lambda_i \in \mathbb{R}$. Notice $\sum_{i=1}^k m_i = \deg(\text{char poly})$

$= n$. Let β_i be a basis for E_{λ_i} . By (b),

know $\#\beta_i = \dim E_{\lambda_i} = m_i$. Set $\beta = \beta_1 \cup \cdots \cup \beta_k$.

Now $\#\beta = \sum \#\beta_i = n$ and β is lin. indep by the lemma. So β is a basis of \mathbb{R}^n .

consisting of eigenvectors for A , and so

A is diagonalizable.

⑤

(\Rightarrow) Suppose A is diagonalizable. Showed last time that (a) follows, so let λ_i, m_i be as before. Set $d_i = \dim E_{\lambda_i}$ which we know satisfies $d_i \leq m_i$. Let β be a basis for \mathbb{R}^n consisting of eigenvectors of A . Set $b_i = \#$ of v in β that are in E_{λ_i} . Know $b_i \leq d_i \leq m_i$ and so

$$n = \# \beta = \sum b_i \leq \sum d_i \leq \sum m_i = n$$

Thus we must have $b_i = d_i = m_i$ for all i , proving (b). Q.E.D.